

Neutrino masses and the number of neutrino species from WMAP and 2dFGRS

Steen Hannestad*

*Department of Physics, University of Southern Denmark,
Campusvej 55, DK-5230 Odense M, Denmark*

and

*NORDITA, Blegdamsvej 17,
DK-2100 Copenhagen, Denmark*

(Dated: February 2, 2008)

We have performed a thorough analysis of the constraints which can be put on neutrino parameters from cosmological observations, most notably those from the WMAP satellite and the 2dF galaxy survey. For this data we find an upper limit on the sum of active neutrino mass eigenstates of $\sum m_\nu \leq 1.0$ eV (95% conf.), but this limit is dependent on priors. We find that the WMAP and 2dF data alone cannot rule out the evidence from neutrinoless double beta decay reported by the Heidelberg-Moscow experiment. In terms of the relativistic energy density in neutrinos or other weakly interacting species we find, in units of the equivalent number of neutrino species, N_ν , that $N_\nu = 4.0^{+3.0}_{-2.1}$ (95 % conf.). When BBN constraints are added, the bound on N_ν is $2.6^{+0.4}_{-0.3}$ (95 % conf.), suggesting that N_ν could possibly be lower than the standard model value of 3. This can for instance be the case in models with very low reheating temperature and incomplete neutrino thermalization. Conversely, if N_ν is fixed to 3 then the data from WMAP and 2dFGRS predicts that $0.2458 \leq Y_P \leq 0.2471$ (95% conf.), which is significantly higher than the observationally measured value. The limit on relativistic energy density changes when a small ν_e chemical potential is present during BBN. In this case the upper bound on N_ν from WMAP, 2dFGRS and BBN is $N_\nu \leq 6.5$. Finally, we find that a non-zero $\sum m_\nu$ can be compensated by an increase in N_ν . One result of this is that the LSND result is not yet ruled out by cosmological observations.

PACS numbers: 14.60.Pq, 95.35.+d, 98.80.-k

I. INTRODUCTION

Neutrinos exist in equilibrium with the electromagnetic plasma in the early universe, until a temperature of a few MeV. At this point the weak interactions freeze out and neutrinos decouple from the plasma. Shortly after this event, the temperature of the plasma falls below the electron mass, and electrons and positrons annihilate, dumping their entropy into the photon gas. This heats the photon gas while having no effect on neutrinos, and the end result is that the photon temperature is larger than the neutrino temperature by the factor $(11/4)^{1/3} \simeq 1.40$. Since the present day photon temperature has been measured with great accuracy to be 2.728 K, the neutrino temperature is known to be 1.95 K, or about 2×10^{-4} eV. Since the heaviest neutrino has a mass of at least about 0.04 eV it must at present be extremely non-relativistic and therefore acts as dark matter. The contribution of a single neutrino species of mass m_ν to the present day matter density can be written as [25, 26, 27]

$$\Omega_\nu h^2 = \frac{m_\nu}{92.5 \text{ eV}}, \quad (1)$$

so that for a neutrino mass of about 30 eV, neutrinos will make up all of the dark matter. However, this would have disastrous consequences for structure for-

mation in the universe, because neutrinos of eV mass have very large free streaming lengths and would erase structure in the neutrino density on scales smaller than $l_{\text{fs}} \simeq 1 m_{\nu, \text{eV}}^{-1}$ Gpc completely. This leads to an overall suppression of matter fluctuations at small scales, an effect which is potentially observable.

A. Absolute value of neutrino masses

The absolute value of neutrino masses are very difficult to measure experimentally. On the other hand, mass differences between neutrino mass eigenstates, (m_1, m_2, m_3) , can be measured in neutrino oscillation experiments. Observations of atmospheric neutrinos suggest a squared mass difference of $\delta m^2 \simeq 3 \times 10^{-3} \text{ eV}^2$ [1, 2, 3]. While there are still several viable solutions to the solar neutrino problem from solar neutrino observations alone, the large mixing angle (LMA) solution gives by far the best fit with $\delta m^2 \simeq 5 \times 10^{-5} \text{ eV}^2$ [4, 5]. Recently the KamLAND reactor neutrino experiment has announced a positive detection of neutrino oscillations indicating that the LMA solution is indeed correct [24].

In the simplest case where neutrino masses are hierarchical these results suggest that $m_1 \sim 0$, $m_2 \sim \delta m_{\text{solar}}$, and $m_3 \sim \delta m_{\text{atmospheric}}$. If the hierarchy is inverted [6, 7, 8, 9, 10, 11] one instead finds $m_3 \sim 0$, $m_2 \sim \delta m_{\text{atmospheric}}$, and $m_1 \sim \delta m_{\text{atmospheric}}$. However, it is also possible that neutrino masses are degenerate [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22], $m_1 \sim m_2 \sim m_3 \gg \delta m_{\text{atmospheric}}$, in which case oscillation

*Electronic address: steen@nordita.dk

experiments are not useful for determining the absolute mass scale.

Experiments which rely on kinematical effects of the neutrino mass offer the strongest probe of this overall mass scale. Tritium decay measurements have been able to put an upper limit on the electron neutrino mass of 2.2 eV (95% conf.) [23]. However, cosmology at present yields an even stronger limit which is also based on the kinematics of neutrino mass. As discussed before any structure in the neutrino density below the free-streaming scale is erased and therefore the presence of a non-zero neutrino mass suppresses the matter power spectrum at small scales relative to large scale, roughly by $\Delta P/P \sim -8\Omega_\nu/\Omega_m$ [75].

This power spectrum suppression allows for a determination of the neutrino mass from measurements of the matter power spectrum on large scales, as well as the spectrum of CMB fluctuations. This matter spectrum is related to the galaxy correlation spectrum measured in large scale structure (LSS) surveys via the bias parameter, $b^2(k) \equiv P_g(k)/P_m(k)$. Such analyses have been performed several times before [78, 79], most recently using data from the 2dFGRS galaxy survey [48, 80, 81]. These investigations found mass limits of 1.5-3 eV, depending on assumptions about the cosmological parameter space.

In a seminal paper it was calculated by Eisenstein, Hu and Tegmark that future CMB and LSS experiments could push the bound on the sum of neutrino masses down to about 0.3 eV [75]. The prospects for an absolute neutrino mass determination was discussed in further detail in Ref. [29] where it was found that in fact the upper bound could be pushed to 0.12 eV (95% conf.) using data from the Sloan Digital Sky Survey and the upcoming Planck satellite.

More recently the new WMAP data, in conjunction with large scale structure data from 2dFGRS has been used to put an upper bound on the sum of all neutrino species of $\sum m_\nu \leq 0.7$ eV (95% conf.) [28].

However, the exact value of this upper bound depends strongly on priors on other cosmological parameters, mainly H_0 . In the present paper we calculate the upper bound on $\sum m_\nu$ from present cosmological data, with an emphasis on studying how the bound depends on the data set chosen.

In addition to their contribution to the cosmological mass density neutrinos also contribute to the cosmological energy density around the epoch of recombination. Neutrinos which have mass smaller than roughly $3T_{\text{rec}}$, where $T_{\text{rec}} \simeq 0.3$ eV is the temperature of recombination, will act as fully relativistic particles when it comes to CMB and large scale structure.

In the standard model there are three light neutrino species with this property. However, these particles are not necessarily in an equilibrium Fermi-Dirac distribution with zero chemical potential. It is well known that the universe contains a non-zero baryon asymmetry of the order $\eta = \frac{n_B - n_{\bar{B}}}{n_\gamma} \sim 10^{-10}$. A neutrino asymmetry of similar magnitude would have no impact on cosmology

during CMB and LSS formation, but since the neutrino asymmetry is not directly observable it could in principle be much larger than the baryon asymmetry. Such a neutrino asymmetry would effectively show up as extra relativistic energy density in the CMB and LSS power spectra.

Another possibility for extra relativistic energy is that there are additional light species beyond the standard model which have decoupled early (such as the graviton or the gravitino).

From the perspective of late time evolution at $T \lesssim 1$ MeV it is customary to parametrize any such additional energy density in terms of N_ν [30], the equivalent number of neutrino species. In Section III we discuss bounds on N_ν from the present WMAP and 2dFGRS data, combined with additional information on other cosmological parameters from the Hubble HST key project and the Supernova Cosmology Project.

However, as will be discussed later, a non-zero neutrino chemical potential can have an effect on big bang nucleosynthesis which is profoundly different from simple relativistic energy density if it is located in the electron neutrino sector. This possibility will be further discussed in Section III.

Another important point is that any entropy production which takes place after BBN, but prior to CMB formation will only be detectable via CMB and LSS observations. One such example is the decay of a hypothetical long-lived massive particle at temperatures below roughly 0.01 MeV.

The general outline of the paper is as follows: In Section II we discuss the likelihood analysis as well as the data sets used in the analysis. In Section III we present the numerical results of the analysis in terms of an upper bound on $\sum m_\nu$, a confidence interval on N_ν , and a bound on the neutrino chemical potentials. Finally, section IV contains a discussion and conclusion.

II. LIKELIHOOD ANALYSIS

The extraction of cosmological parameters from cosmological data is a difficult process since for both CMB and LSS the power spectra depend on a plethora of different parameters. Furthermore, since the CMB and matter power spectra depend on many different parameters one might worry that an analysis which is too restricted in parameter space could give spuriously strong limits on a given parameter. We discuss this point in further detail below.

For calculating the theoretical CMB and matter power spectra we use the publicly available CMBFAST package [31]. As the set of cosmological parameters we choose Ω_m , the matter density, the curvature parameter, Ω_b , the baryon density, H_0 , the Hubble parameter, n_s , the scalar spectral index of the primordial fluctuation spectrum, τ , the optical depth to reionization, Q , the normalization of the CMB power spectrum, b , the bias parameter, and

finally the two parameters related to neutrino physics, $\Omega_\nu h^2$ and N_ν . We restrict the analysis to geometrically flat models, i.e. $\Omega = \Omega_m + \Omega_\Lambda = 1$.

In principle one might include even more parameters in the analysis, such as r , the tensor to scalar ratio of primordial fluctuations. However, r is most likely so close to zero that only future high precision experiments may be able to measure it. The same is true for other additional parameters. Small deviations from slow-roll during inflation can show up as a logarithmic correction to the simple power-law spectrum predicted by slow-roll. [32, 33, 34] or additional relativistic energy density [35, 36, 37, 38, 39, 40, 41, 42, 43, 44] could be present. However, there is no evidence of any such effect in the present data and therefore we restrict the analysis to the “minimal” standard cosmological model plus neutrino physics.

In this full numerical likelihood analysis we use the free parameters discussed above with certain priors determined from cosmological observations other than CMB and LSS. In flat models the matter density is restricted by observations of Type Ia supernovae to be $\Omega_m = 0.28 \pm 0.14$ [45], and the HST Hubble key project has obtained a limit on H_0 of $72 \pm 8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [46].

We calculate bounds on neutrino physics, both with and without, the constraints on Ω_m and H_0 . In the cases where we do not use the SNI-a and HST priors we use simple top-hat priors instead, $0.1 \leq \Omega_m \leq 0.5$ and $0.5 \leq h \leq 0.85$.

Particularly the prior on H_0 is of great importance in constraining neutrino parameter space.

The numerical marginalization over parameters other than $\Omega_\nu h^2$ was performed using a simulated annealing procedure [47]. A full description of how χ^2 is calculated can for instance be found in [48].

A. LSS data

At present, by far the largest survey available is the 2dFGRS [49] of which about 147,000 galaxies have so far been analyzed. Tegmark, Hamilton and Xu [50] have calculated a power spectrum, $P(k)$, from this data, which we use in the present work. The 2dFGRS data extends to very small scales where there are large effects of non-linearity. Since we only calculate linear power spectra, we use (in accordance with standard procedure) only data on scales larger than $k = 0.2h \text{ Mpc}^{-1}$, where effects of non-linearity should be minimal [29]. Making this cut reduces the number of power spectrum data points to 18.

B. k -dependent bias

In all present parameter estimation analyses it is assumed that the bias parameter, $b^2(k) \equiv P_g(k)/P_m(k)$, is independent of the scale, k .

However, many independent simulations find that bias is in fact quite strongly scale dependent [51, 52] in the non-linear regime. In the linear regime bias is expected to be constant, and the asymptotic value $b_\infty = \lim_{k \rightarrow 0} b(k)$ is reached as a scale of roughly $k \simeq 0.1 - 0.2h \text{ Mpc}^{-1}$. This means that at scales larger than k_{cut} bias should be very close to scale-independent, and that we can therefore use a single parameter, b , to describe it.

C. CMB data

The CMB temperature fluctuations are conveniently described in terms of the spherical harmonics power spectrum

$$C_l \equiv \langle |a_{lm}|^2 \rangle, \quad (2)$$

where

$$\frac{\Delta T}{T}(\theta, \phi) = \sum_{lm} a_{lm} Y_{lm}(\theta, \phi). \quad (3)$$

Since Thomson scattering polarizes light there are additional powerspectra coming from the polarization anisotropies. The polarization can be divided into a curl-free (E) and a curl (B) component, yielding four independent power spectra: $C_{T,l}$, $C_{E,l}$, $C_{B,l}$ and the temperature E -polarization cross-correlation $C_{TE,l}$.

The WMAP experiment have reported data only on $C_{T,l}$ and $C_{TE,l}$, as described in Ref. [28, 53, 54, 55, 56]

We have performed the likelihood analysis using the prescription given by the WMAP collaboration which includes the correlation between different C_l 's [28, 53, 54, 55, 56]. Foreground contamination has already been subtracted from their published data.

In parts of the data analysis we also add other CMB data from the compilation by Wang *et al.* [74] which includes data at high l . Altogether this data set has 28 data points.

III. NUMERICAL RESULTS

A. Neutrino masses

We have calculated χ^2 as a function of neutrino mass while marginalizing over all other cosmological parameters. This has been done using the data sets described above. In the first case we have calculated the constraint using the WMAP $C_{T,l}$ combined with the 2dFGRS data, and in the second case we have added the polarization measurement from WMAP. Finally we have added the additional constraint from the HST key project and the Supernova Cosmology Project. It should also be noted that when constraining the neutrino mass it has in all cases been assumed that N_ν is equal to the standard model value of 3.04. In section III.D we relax this condition in order to study the LSND bound.

The result is shown in Fig. 1. As can be seen from the figure the 95% confidence upper limit on the sum of neutrino masses is $\sum m_\nu \leq 1.01$ eV (95% conf.) using the case with priors. This value is completely consistent with what is found in Ref. [82] where simple Gaussian priors from WMAP were added to the 2dFGRS data analysis. For the three cases studied we find the following limits:

$$\begin{aligned} \sum m_\nu &< 1.01 \text{ eV} && \text{for WMAP+Wang+2dFGRS+HST+SN-Ia} \\ \sum m_\nu &< 1.20 \text{ eV} && \text{for WMAP+Wang+2dFGRS} \\ \sum m_\nu &< 2.12 \text{ eV} && \text{for WMAP+2dFGRS} \end{aligned}$$

However, it is somewhat higher than the upper limit of $\sum m_\nu \leq 0.7$ eV found in the WMAP analysis [28]. There are several reasons for this: First, we do not use Ly- α forest data in our analysis. While adding them to the analysis clearly would improve the limit the interpretation of Ly- α data is still somewhat controversial [84]. Therefore the limit derived here can be viewed as more robust, albeit also more conservative. The second reason is that we use a completely free bias parameter. It well known that a determination of the neutrino mass is to some extent degenerate with the bias parameter b so that a precise determination of b would shrink the allowed range even further. However, the bias parameter is probably the most poorly understood parameter in the analysis of large scale structure [74] and therefore imposing a particular prior on b could yield too restrictive results. This is the reason why we do not show any result for χ^2 with a b -prior.

Also, for accurate CMB and LSS data sets, the main degeneracy is not with the bias parameter, but rather with the Hubble parameter. This was also noted in Ref. [29].

In the middle panel of Fig. 1 we show the best fit value of H_0 for a given $\Omega_\nu h^2$. It is clear that an increasing value of $\sum m_\nu$ can be compensated by a decrease in H_0 . Even though the data yields a strong constraint on $\Omega_m h^2$ there is no independent constraint on Ω_m in itself. Therefore, an decreasing H_0 leads to an increasing Ω_m . This can be seen in the bottom panel of Fig. 1.

When the HST prior on H_0 is relaxed a higher value of $\sum m_\nu$ is allowed, in the case with only WMAP and 2dFGRS data the upper bound is $\Omega_\nu h^2 \leq 0.023$ (95% conf.), corresponding to a neutrino mass of 0.71 eV for each of the three neutrinos.

This effect was also found by Elgarøy and Lahav [82] in their analysis of the effects of priors on the determination of $\sum m_\nu$.

However, as can also be seen from the figure, the addition of high- l CMB data from the Want *et al.* compilation also shrinks the allowed range of $\sum m_\nu$ significantly. The reason is that there is a significant overlap of the scales probed by high- l CMB experiments and the 2dFGRS survey. Therefore, even though we use bias as a free fitting parameter, it is strongly constrained by the fact that the CMB and 2dFGRS data essentially cover much of the same range in k -space.




FIG. 1: The top panel shows χ^2 as a function of $\sum m_\nu$ for different choices of priors. The dotted line is for WMAP + 2dFGRS data alone, the dashed line is with the additional Wang *et al.* data. The full line is for additional HST and SNI-a priors as discussed in the text. The horizontal lines show $\Delta\chi^2 = 1$ and 4 respectively. The middle panel shows the best fit values of H_0 for a given $\sum m_\nu$. The horizontal lines show the HST key project 1σ limit of $H_0 = 72 \pm 8$ km s Mpc $^{-1}$. Finally, the lower panel shows best fit values of Ω_m . In this case the horizontal line corresponds to the SNI-a 1σ upper limit of $\Omega_m < 0.42$.

It should be noted that Elgarøy and Lahav [82] find that bias does not play any role in determining the bound on $\sum m_\nu$. At first this seems to contradict the discussion here, and also what was found from a Fisher matrix analysis in Ref. [48]. The reason is that in Ref. [82], redshift distortions are included in the 2dFGRS data analysis. Given a constraint on the amplitude of fluctuations from CMB data, and a constraint on $\Omega_m h^2$, this effectively constrains the bias parameter. Therefore adding a further constraint on bias in their analysis does not change the results.

Neutrinoless double beta decay

Recently it was claimed that the Heidelberg-Moscow experiment yields positive evidence for neutrinoless double beta decay. Such experiments probe the ‘effective electron neutrino mass $m_{ee} = |\sum_j U_{ej}^2 m_{\nu_j}|$. Given the

uncertainties in the involved nuclear matrix elements the Heidelberg-Moscow result leads to a mass of $m_{ee} = 0.3 - 1.4$ eV. If this is true then the mass eigenstates are necessarily degenerate, and $\sum m_\nu \simeq 3m_{ee}$. Taking the WMAP result of $\sum m_\nu \leq 0.70$ eV at face value seems to be inconsistent with the Heidelberg-Moscow result [85]. However, already if Ly- α forest data and a constraint on the bias parameter is not used in the analysis the upper bound of $\sum m_\nu \leq 1.01$ eV is still consistent. For this reason it is probably premature to rule out the claimed evidence for neutrinoless double beta decay.

B. Neutrino relativistic energy density

For the case of the effective number of neutrino species we have in all cases calculated constraints in the $(\Omega_b h^2, N_\nu)$ plane, while marginalizing over all other parameters. The reason for this is that for Big Bang Nucleosynthesis purposes these are exactly the important parameters. Therefore, to combine CMB, LSS, and BBN constraints the marginalization over $\Omega_b h^2$ should not be performed. Furthermore, when constraining N_ν we have always assumed that $\sum m_\nu \simeq 0$ so that the neutrino mass has no influence on cosmology.

We start out by investigating constraints on N_ν from CMB and LSS data alone. In Fig. 2 we show $\Delta\chi^2$ for a global fit to N_ν which marginalizes over all other cosmological parameters. The overall best fit for the WMAP T and TE data, combined with the Wang *et al.* compilation, the 2dFGRS data, the HST key project data on H_0 , as well as the SNI-a data on Ω_m , has $\chi^2 = 1467.6$ for a total of 1395 degrees of freedom. This gives $\chi^2/\text{d.o.f.} = 1.05$ which is entirely compatible with the best fit WMAP value for the standard Λ CDM model of $\chi^2/\text{d.o.f.} = 1.066$. We also show constraints for two other analyses. The first is for WMAP and 2dFGRS data alone and the second for WMAP data alone. The bounds for the three cases are

$$\begin{aligned} N_\nu &= 4.0^{+3.0}_{-2.1} && \text{for WMAP+Wang+2dFGRS+HST+SN-Ia} \\ N_\nu &= 3.1^{+3.9}_{-2.8} && \text{for WMAP+2dFGRS} \\ N_\nu &= 2.1^{+6.7}_{-2.2} && \text{for WMAP only} \end{aligned}$$

These bound are entirely compatible with those found by Crotty, Lesgourgues and Pastor [73], and much tighter than the pre-WMAP constraints.

The constraints derived here are also compatible with what is found by Pierpaoli [72], where an assumption of spatial flatness was relaxed.

In the lower panels of Fig. 2 we show the best fit values of H_0 and Ω_m for a given value of N_ν . The main point to note is that the constraint on N_ν is strongly dependent on H_0 . This was also found in Ref. [40]. With only CMB data and the weak top-hat prior on H_0 the bound on N_ν is very weak. Adding the HST Key Project prior on H_0 cut away a significant amount of parameter space both at low and high N_ν . Adding the 2dFGRS and Wang *et al.* data mainly has the effect of shifting the best fit value




FIG. 2: χ^2 as a function of $\sum N_\nu$ for different choices of priors. The dotted line is for WMAP data alone, the dashed line is with the additional Wang *et al.* and 2dFGRS data. The full line is for additional HST and SNI-a priors as discussed in the text. The horizontal lines show $\Delta\chi^2 = 1$ and 4 respectively. The middle panel shows the best fit values of H_0 for a given N_ν . The horizontal lines show the HST key project 1σ limit of $H_0 = 72 \pm 8$ km s Mpc $^{-1}$. Finally, the lower panel shows best fit values of Ω_m . In this case the horizontal line corresponds to the SNI-a 1σ upper limit of $\Omega_m < 0.42$.

to higher N_ν , but also cuts away the low N_ν values, an effect also seen in Ref. [73].

In Fig. 3 we show constraints on $(\Omega_b h^2, N_\nu)$ for the full data set described above. The best fit value for $\Omega_b h^2$ is 0.0233, which is equivalent to the value found in the WMAP data analysis. In the 2-dimensional plots the 68% and 95% regions are formally defined by $\Delta\chi^2 = 2.30$ and 6.17 respectively. Note that this means that the 68% and 95% contours are not necessarily equivalent to the same confidence level for single parameter estimates.

It should be noted here that in addition to an upper bound on N_ν there is also a 3.0σ confidence detection of $N_\nu > 0$. This is in concordance with the pre-WMAP data from which a non-trivial lower bound on N_ν could also be derived.

Adding BBN information – In the case where all the relativistic energy density contained in N_ν is produced prior to BBN, a BBN constraint can be added to the




FIG. 3: 68% and 95% confidence contours in the $(\Omega_b h^2, N_\nu)$ plane for the WMAP TT and TE data, combined with the 2dFGRS data, the HST data on H_0 and the SNI-a data on Ω_m .

CMB and LSS constraint without any problems. In practice we have used abundances of He-4 and D to make constraints in the $(\Omega_b h^2, N_\nu)$ plane. We use the following values for the primordial abundances [63, 64]

$$D/H = 2.78^{+0.44}_{-0.38} \times 10^{-5} \quad (4)$$

$$Y_P = 0.238 \pm 0.005 \quad (5)$$

This calculation is shown in Fig. 4. In terms of a single parameter constraint on N_ν it is $N_\nu = 2.6^{+0.4}_{-0.3}$ (95 % conf.). Compared to the recent calculation by Abazajian [62] of a BBN-only constraint of $1.7 \leq N_\nu \leq 3.5$ (95 % conf.) this is a significant improvement. Very interestingly the new limit suggests the possibility that N_ν is actually less than 3. This is for instance possible in scenarios with extremely low reheating temperature [65, 66, 67].

Of course this conclusion is mainly based on the fact that CMB and LSS data prefers a slightly higher value of $\Omega_b h^2$ than BBN. It should also be stressed that the estimates of the primordial abundances could be biased by systematic effects so that the quoted statistical error bar is not really meaningful. Therefore it is probably premature to talk of any inconsistency between the $N_\nu = 3$ prediction of the standard model and observations.

In fact the argument can also be reversed. If N_ν is fixed to the standard model value of 3 then then CMB and LSS constraint on $\Omega_b h^2$ provides an accurate measurement of primordial He-4. Using the derived constraint on $\Omega_b h^2$ the 95% confidence range for Y_P is

$$0.2458 \leq Y_P \leq 0.2471. \quad (6)$$

This could point to a serious underlying systematic effect




FIG. 4: 68% and 95% confidence contours in the $(\Omega_b h^2, N_\nu)$ plane for the same data sets as in fig. 3, but with the addition of BBN data. The lined contours are the 68% and 95% regions for BBN data alone.

in observational determinations of Y_P , as discussed in Ref. [92].

Late entropy production – However, if entropy is produced subsequent to BBN and prior to CMB formation it still behaves like an additional N_ν for CMB and LSS data, while having no effect on BBN. In the case where the decay of a massive particle produces the entropy the bound on N_ν can be translated into a bound on the mass and lifetime of the massive particle. The effective N_ν produced by complete decay of a non-relativistic species at $10\text{keV} \gtrsim T \gtrsim 1\text{ eV}$ is

$$N_{\nu,\text{eff}} \simeq 3 + 1.2 \times \left(\frac{n}{n_{\nu,0}} \right) m_{\text{keV}}^{4/3} \tau_y^{2/3}, \quad (7)$$

where $n_{\nu,0}$ is the number density of a standard model neutrino.

The bound on N_ν therefore translates into a bound of

$$\left(\frac{n}{n_{\nu,0}} \right) m_{\text{keV}}^{4/3} \tau_y^{2/3} \leq 3.3 \quad (8)$$

for any hypothetical massive particle.

C. Neutrino lepton asymmetry and N_ν

Whether the extra relativistic energy density is in the form of a non-zero chemical potential has no influence on CMB or LSS calculations since they are not sensitive to the flavour content of the additional energy density. However, from a BBN perspective neutrinos in the electron flavour are very different from muon and tau neutrinos. The reason is that electron neutrinos and antineutrinos

enter directly into the weak interactions which maintain neutron-proton equilibrium. Adding electron neutrinos shifts the balance in the direction of a lower n/p ratio, i.e. the opposite effect of increasing N_ν . Therefore a large chemical potential in the muon or tau neutrino flavours can be hidden by compensating its effect on BBN with a small electron neutrino chemical potential. In fact, the additional N_ν would not necessarily have to be in the form of a non-zero ν_μ or ν_τ chemical potential. A large N_ν from additional weakly interacting species could also be compensated by a small electron neutrino chemical potential.

However, since CMB and LSS observations are insensitive to flavour this degeneracy can be broken by combining BBN, CMB and SNI-a data [41]. The effect of doing this can be seen in Fig. 5. From the present data a bound on the chemical potential in non-electron flavour neutrinos is $|\eta| \leq 2.3$ at (95% conf.), where $\eta = \mu/T$. In terms of N_ν the bound is $N_\nu \leq 6.5$. As can be seen from the figure this bound comes almost entirely from the CMB+LSS data alone.

It should be noted that any degree of neutrino mixing will act to equilibrate the flavours and possibly eliminate this masking effect. For small mass differences neutrino oscillations occur after weak freeze-out and consequently there is no effect on BBN. However, given the mass differences and close to maximal mixing angles suggested by the solar and atmospheric neutrino data, all three flavours will be almost equilibrated before weak freeze-out. This means that the bound on the lepton asymmetry in all flavours will be close to the BBN bound for the electron sector, i.e. $|\eta_e| \lesssim 0.15$ [44, 68, 69, 70, 71]. Therefore the bound derived here is not competitive, but is of course very robust because it does not depend on a knowledge of mass differences or mixing angles. It is also more robust in the sense that a large N_ν in hypothetical new particles can still be invisible to BBN if all neutrino species have a small chemical potential. Therefore the N_ν bound derived in this section does not have to be in the form of a $\nu_{\mu,\tau}$ chemical potential.

D. Combining $\sum m_\nu$ with N_ν - constraining LSND

From the analyses in the above two sections it was found that: (a) An increasing $\sum m_\nu$ can be compensated by a decreasing H_0 and (b) An increasing N_ν can be compensated by an increasing H_0 . One might therefore wonder whether a model with non-zero $\sum m_\nu$, combined with $N_\nu > 3$ can provide a good fit to the data.

In order to test this we have performed a likelihood analysis for $\sum m_\nu$ for different values of N_ν . We show this in Fig. 6. This analysis was performed with all available data and priors.

As can be seen from the figure, the best fit actually is actually shifted to higher $\sum m_\nu$ when N_ν increases, and the conclusion is that a model with high neutrino mass and additional relativistic energy density can provide ac-




FIG. 5: The full lines show 68% and 95% confidence regions in the $(\Omega_b h^2, N_\nu)$ plane for the case where the additional N_ν is compensated during BBN by a small ν_e chemical potential. The full contours are for BBN data alone, whereas the dashed lines are for CMB and LSS data.




FIG. 6: $\Delta\chi^2$ as a function of $\sum m_\nu$ for various different values of N_ν . The full line is for $N_\nu = 3$, the dotted for $N_\nu = 4$, and the dashed for $N_\nu = 5$. $\Delta\chi^2$ is calculated relative to the best fit $N_\nu = 3$ model.

ceptable fits to the data. As a function of N_ν the upper bound on $\sum m_\nu$ is (at 95% confidence)

$$\begin{aligned} \sum m_\nu &\leq 1.01 \text{ eV} & \text{for } N_\nu = 3 \\ \sum m_\nu &\leq 1.38 \text{ eV} & \text{for } N_\nu = 4 \\ \sum m_\nu &\leq 2.12 \text{ eV} & \text{for } N_\nu = 5 \end{aligned}$$

This has significant implications for attempts to constrain the LSND experiment using the present cosmological data. Pierce and Murayama conclude from the

present MAP limit that the LSND result is excluded [85] (see also Ref. [86]).

However, for several reasons this conclusion does not follow trivially from the present data. In general the three mass differences implied by Solar, atmospheric and the LSND neutrino measurements can be arranged into either 2+2 or 3+1 schemes. Recent analyses [93] of experimental data have shown that the 2+2 models are ruled out. The 3+1 scheme with a single massive state, m_4 , which makes up the LSND mass gap, is still marginally allowed in a few small windows in the $(\Delta m^2, \sin^2 2\theta)$ plane. These gaps are at $(\Delta m^2, \sin^2 2\theta) \simeq (0.8 \text{ eV}^2, 2 \times 10^{-3})$, $(1.8 \text{ eV}^2, 8 \times 10^{-4})$, $(6 \text{ eV}^2, 1.5 \times 10^{-3})$ and $(10 \text{ eV}^2, 1.5 \times 10^{-3})$. These four windows corresponds to masses of 0.9, 1.4, 2.5 and 3.2 eV respectively. From the Solar and atmospheric neutrino results the three light mass eigenstates contribute only about 0.1 eV of mass if they are hierarchical. This means that the sum of all mass eigenstate is close to m_4 .

The limit for $N_\nu = 4$ which corresponds roughly to the LSND scenario is $\sum m_\nu \lesssim 1.4 \text{ eV}$, which still leaves the lowest of the remaining windows. The second window at $m \sim 1.8 \text{ eV}$ is disfavoured by the data, but not at very high significance.

The second reason why the LSND sterile neutrino is not necessarily ruled out is that it was not necessarily fully equilibrated in the early universe. The cosmological mass limit applies only to a species with decoupling temperature around 1 MeV. If a sterile species is not fully equilibrated its number density is lower than that of a standard active neutrino species and therefore the mass limit will be less restrictive. In the simplest possible case a neutrino in a the lowest of the four LSND windows will have an abundance of roughly 0.5 times that of a standard active neutrino, whereas neutrino masses and mixings corresponding to the other three windows will lead to almost perfect equilibration [87, 88]. However, for instance the presence of a non-zero lepton number can strongly suppress the production of sterile neutrinos [89, 90].

IV. DISCUSSION

We have calculated improved constraints on neutrino masses and the cosmological relativistic energy density, using the new WMAP data together with data from the 2dFGRS galaxy survey.

Using CMB and LSS data together with a prior from the HST key project on H_0 yielded an upper bound of $\sum m_\nu \leq 1.01 \text{ eV}$ (95% conf.). While this excludes most

of the parameter range suggested by the claimed evidence for neutrinoless double beta decay in the Heidelberg-Moscow experiment, it seems premature to rule out this claim based on cosmological observations.

Another issue where the cosmological upper bound on neutrino masses is very important is for the prospects of directly measuring neutrino masses in tritium endpoint measurements. The successor to the Mainz experiment, KATRIN, is designed to measure an electron neutrino mass of roughly 0.25 eV [91], or in terms of the sum of neutrino mass eigenstates, $\sum m_\nu \lesssim 0.75 \text{ eV}$. The WMAP result of $\sum m_\nu \lesssim 0.7 \text{ eV}$ (95% conf.) already seems to exclude a positive measurement of mass in KATRIN. However, this very tight limit depends on priors, as well as Ly- α forest data, and the more conservative present limit of $\sum m_\nu \lesssim 1.01 \text{ eV}$ (95% conf.) does not exclude that KATRIN will detect a neutrino mass.

From the data we also inferred a limit on N_ν of $N_\nu = 4.0_{-2.1}^{+3.0}$ (95% conf.) on the equivalent number of neutrino species. This is a marked improvement over the previous best limit of roughly $N_\nu \leq 13$ [40].

When light element measurements of He-4 and D are included the bound is strengthened considerably to $N_\nu = 2.6_{-0.3}^{+0.4}$ (95 % conf.). Interestingly this suggests a possible value of N_ν which is *less* than 3. This could be the case for instance in scenarios with very low reheating temperature where neutrinos were never fully equilibrated [65, 66, 67]. However, it should be stressed that primordial abundances could be dominated by systematics. Therefore it is probably premature to talk of a new BBN “crisis”.

The bound on N_ν also translates into a bound on a possible neutrino lepton asymmetry in non-electron neutrinos of $|\eta_\nu| \leq 2.4$. Even though this bound is much stronger than the previous bound of 2.6 [41] it is weak compared to the bound obtainable from BBN considerations alone when flavour oscillations are accounted for [44, 68, 69, 70, 71].

Finally, we also found that the neutrino mass bound depends on the total number of light neutrino species. In scenarios with sterile neutrinos this is an important factor. For instance in 3+1 models the mass bound increases from 1.0 eV to 1.4 eV, meaning that the LSND result is not ruled out by cosmological observations yet.

Acknowledgments

I wish to thank Øystein Elgarøy, Julien Lesgourgues, and Sergio Pastor for extensive and enlightening discussions

-
- [1] N. Fornengo, M. C. Gonzalez-Garcia and J. W. Valle, Nucl. Phys. B **580** (2000) 58 [arXiv:hep-ph/0002147].
 [2] S. Fukuda *et al.* [Super-Kamiokande Collaboration],

- Phys. Rev. Lett. **85**, 3999 (2000) [arXiv:hep-ex/0009001].
 [3] M. Maltoni, T. Schwetz, M. A. Tortola and J. W. Valle, arXiv:hep-ph/0207227.

- [4] J. N. Bahcall, M. C. Gonzalez-Garcia and C. Pena-Garay, arXiv:hep-ph/0204314.
- [5] See the SNO homepage <http://sno.phy.queensu.ca>
- [6] V. A. Kostelecky and S. Samuel, Phys. Lett. B **318**, 127 (1993).
- [7] G. M. Fuller, J. R. Primack and Y. Z. Qian, Phys. Rev. D **52**, 1288 (1995) [arXiv:astro-ph/9502081].
- [8] D. O. Caldwell and R. N. Mohapatra, Phys. Lett. B **354**, 371 (1995) [arXiv:hep-ph/9503316].
- [9] S. M. Bilenky, C. Giunti, C. W. Kim and S. T. Petcov, Phys. Rev. D **54**, 4432 (1996) [arXiv:hep-ph/9604364].
- [10] S. F. King and N. N. Singh, Nucl. Phys. B **596**, 81 (2001) [arXiv:hep-ph/0007243].
- [11] H. J. He, D. A. Dicus and J. N. Ng, arXiv:hep-ph/0203237.
- [12] A. Ioannian and J. W. Valle, Phys. Lett. B **332**, 93 (1994) [arXiv:hep-ph/9402333].
- [13] P. Bamert and C. P. Burgess, Phys. Lett. B **329**, 289 (1994) [arXiv:hep-ph/9402229].
- [14] R. N. Mohapatra and S. Nussinov, Phys. Lett. B **346**, 75 (1995) [arXiv:hep-ph/9411274].
- [15] H. Minakata and O. Yasuda, Phys. Rev. D **56**, 1692 (1997) [arXiv:hep-ph/9609276].
- [16] F. Vissani, arXiv:hep-ph/9708483.
- [17] H. Minakata and O. Yasuda, Nucl. Phys. B **523**, 597 (1998) [arXiv:hep-ph/9712291].
- [18] J. R. Ellis and S. Lola, Phys. Lett. B **458**, 310 (1999) [arXiv:hep-ph/9904279].
- [19] J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. B **556**, 3 (1999) [arXiv:hep-ph/9904395].
- [20] J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. B **569**, 82 (2000) [arXiv:hep-ph/9905381].
- [21] E. Ma, J. Phys. G **25**, L97 (1999) [arXiv:hep-ph/9907400].
- [22] R. Adhikari, E. Ma and G. Rajasekaran, Phys. Lett. B **486**, 134 (2000) [arXiv:hep-ph/0004197].
- [23] J. Bonn *et al.*, Nucl. Phys. Proc. Suppl. **91**, 273 (2001).
- [24] K. Eguchi *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **90**, 021802 (2003) [arXiv:hep-ex/0212021].
- [25] S. Hannestad and J. Madsen, Phys. Rev. D **52**, 1764 (1995) [arXiv:astro-ph/9506015].
- [26] A. D. Dolgov, S. H. Hansen and D. V. Semikoz, Nucl. Phys. B **503**, 426 (1997) [arXiv:hep-ph/9703315].
- [27] G. Mangano, G. Miele, S. Pastor and M. Peloso, arXiv:astro-ph/0111408.
- [28] D. N. Spergel *et al.*, astro-ph/0302209
- [29] S. Hannestad, arXiv:astro-ph/0211106.
- [30] G. Steigman, D. N. Schramm and J. E. Gunn, Phys. Lett. B **66**, 202 (1977).
- [31] U. Seljak and M. Zaldarriaga, Astrophys. J. **469**, 437 (1996).
- [32] S. Hannestad, S. H. Hansen and F. L. Villante, Astropart. Phys. **16**, 137 (2001) [arXiv:astro-ph/0012009].
- [33] S. Hannestad, S. H. Hansen, F. L. Villante and A. J. Hamilton, Astropart. Phys. **17**, 375 (2002) [arXiv:astro-ph/0103047].
- [34] L. M. Griffiths, J. Silk and S. Zaroubi, astro-ph/0010571.
- [35] G. Jungman, M. Kamionkowski, A. Kosowsky and D. N. Spergel, Phys. Rev. D **54**, 1332 (1996) [arXiv:astro-ph/9512139].
- [36] J. Lesgourgues and M. Peloso, Phys. Rev. D **62**, 081301 (2000) [arXiv:astro-ph/0004412].
- [37] S. Hannestad, Phys. Rev. Lett. **85**, 4203 (2000) [arXiv:astro-ph/0005018].
- [38] S. Esposito, G. Mangano, A. Melchiorri, G. Miele and O. Pisanti, Phys. Rev. D **63**, 043004 (2001) [arXiv:astro-ph/0007419].
- [39] J. P. Kneller, R. J. Scherrer, G. Steigman and T. P. Walker, Phys. Rev. D **64**, 123506 (2001) [arXiv:astro-ph/0101386].
- [40] S. Hannestad, Phys. Rev. D **64**, 083002 (2001) [arXiv:astro-ph/0105220].
- [41] S. H. Hansen, G. Mangano, A. Melchiorri, G. Miele and O. Pisanti, Phys. Rev. D **65**, 023511 (2002) [arXiv:astro-ph/0105385].
- [42] R. Bowen, S. H. Hansen, A. Melchiorri, J. Silk and R. Trotta, arXiv:astro-ph/0110636.
- [43] J. P. Kneller and G. Steigman, arXiv:astro-ph/0210500.
- [44] A. D. Dolgov, S. H. Hansen, S. Pastor, S. T. Petcov, G. G. Raffelt and D. V. Semikoz, arXiv:hep-ph/0201287.
- [45] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], Astrophys. J. **517**, 565 (1999) [arXiv:astro-ph/9812133].
- [46] W. L. Freedman *et al.*, Astrophys. J. Lett. **553**, 47 (2001).
- [47] S. Hannestad, Phys. Rev. D **61**, 023002 (2000).
- [48] S. Hannestad, arXiv:astro-ph/0205223.
- [49] J. Peacock *et al.*, Nature **410**, 169 (2001).
- [50] M. Tegmark, A. J. S. Hamilton and Y. Xu, arXiv:astro-ph/0111575
- [51] M. Blanton *et al.*, Astrophys. J. **531**, 1 (2000)
- [52] R. G. Mann, J. A. Peacock and A. F. Heavens, Mon. Not. R. Astron. Soc. **293**, 209 (1998).
- [53] C. L. Bennett *et al.*, astro-ph/0302207
- [54] A. Kogut *et al.*, astro-ph/0302213
- [55] G. Hinshaw *et al.*, astro-ph/0302217
- [56] L. Verde, *et al.*, astro-ph/0302218
- [57] C. B. Netterfield *et al.* [Boomerang Collaboration], arXiv:astro-ph/0104460.
- [58] A. T. Lee *et al.*, arXiv:astro-ph/0104459.
- [59] N. W. Halverson *et al.*, arXiv:astro-ph/0104489.
- [60] P. F. Scott *et al.*, astro-ph/0205380
- [61] B. S. Mason *et al.*, astro-ph/0205384
- [62] K. N. Abazajian, arXiv:astro-ph/0205238.
- [63] D. Kirkman *et al.*, astro-ph/0302006
- [64] K. A. Olive, G. Steigman and T. P. Walker, Phys. Rept. **333**, 389 (2000) [arXiv:astro-ph/9905320].
- [65] G. F. Giudice, E. W. Kolb and A. Riotto, Phys. Rev. D **64**, 023508 (2001) [arXiv:hep-ph/0005123].
- [66] M. Kawasaki, K. Kohri and N. Sugiyama, Phys. Rev. D **62**, 023506 (2000) [arXiv:astro-ph/0002127].
- [67] P. Adhya and D. R. Chaudhuri, arXiv:hep-ph/0203142.
- [68] C. Lunardini and A. Y. Smirnov, Phys. Rev. D **64**, 073006 (2001) [arXiv:hep-ph/0012056].
- [69] S. Pastor, G. G. Raffelt and D. V. Semikoz, arXiv:hep-ph/0109035.
- [70] K. N. Abazajian, J. F. Beacom and N. F. Bell, Phys. Rev. D **66**, 013008 (2002) [arXiv:astro-ph/0203442].
- [71] Y. Y. Wong, Phys. Rev. D **66**, 025015 (2002) [arXiv:hep-ph/0203180].
- [72] E. Pierpaoli, arXiv:astro-ph/0302465.
- [73] P. Crotty, J. Lesgourgues and S. Pastor, arXiv:astro-ph/0302337.
- [74] X. Wang, M. Tegmark, B. Jain and M. Zaldarriaga, arXiv:astro-ph/0212417.
- [75] W. Hu, D. J. Eisenstein and M. Tegmark, Phys. Rev. Lett. **80**, 5255 (1998) [arXiv:astro-ph/9712057].
- [76] A. G. Doroshkevich, Y. B. Zeldovich and R. A. Sunyaev,

- Sov. Astron. Lett. **6**, 252 (1980)
- [77] A. G. Doroshkevich, M. Y. Khlopov, R. A. Sunyaev, Y. B. Zeldovich and A. S. Szalay, *In *Baltimore 1980, Proceedings, Relativistic Astrophysics**, 32-42.
 - [78] R. A. Croft, W. Hu and R. Dave, Phys. Rev. Lett. **83**, 1092 (1999) [arXiv:astro-ph/9903335].
 - [79] M. Fukugita, G. C. Liu and N. Sugiyama, Phys. Rev. Lett. **84**, 1082 (2000) [arXiv:hep-ph/9908450].
 - [80] Ø. Elgarøy *et al.*, arXiv:astro-ph/0204152.
 - [81] A. Lewis and S. Bridle, arXiv:astro-ph/0205436.
 - [82] Ø. Elgarøy and O. Lahav, astro-ph/0303089.
 - [83] H. V. Klapdor-Kleingrothaus, A. Dietz, H. L. Harney and I. V. Krivosheina, Mod. Phys. Lett. A **16**, 2409 (2001) [arXiv:hep-ph/0201231];
 - [84] R. A. Croft *et al.*, Astrophys. J. **581**, 20 (2002) [arXiv:astro-ph/0012324]; U. Seljak, P. McDonald and A. Makarov, arXiv:astro-ph/0302571;
 - [85] A. Pierce and H. Murayama, arXiv:hep-ph/0302131.
 - [86] C. Giunti, arXiv:hep-ph/0302173.
 - [87] K. Enqvist, K. Kainulainen and M. J. Thomson, Nucl. Phys. B **373**, 498 (1992).
 - [88] S. Hannestad and G. Raffelt, Phys. Rev. D **59**, 043001 (1999) [arXiv:astro-ph/9805223].
 - [89] R. Foot and R. R. Volkas, nucleosynthesis bounds,” Phys. Rev. D **55**, 5147 (1997) [arXiv:hep-ph/9610229].
 - [90] K. Abazajian, G. M. Fuller and M. Patel, Phys. Rev. D **64**, 023501 (2001) [arXiv:astro-ph/0101524].
 - [91] <http://ik1au1.fzk.de/~katrin/>
 - [92] R. H. Cyburt, B. D. Fields and K. A. Olive, arXiv:astro-ph/0302431.
 - [93] M. Maltoni, T. Schwetz, M. A. Tortola and J. W. Valle, Nucl. Phys. B **643**, 321 (2002) [arXiv:hep-ph/0207157].